



Bobtails Rabbit & Guinea Pig Rescue

44 Arundel Avenue
East Ewell
Surrey
KT17 2RG

Tel: 020 8393 2016
Fax: 020 8393 5390

bobtailsrescue@aol.com
www.bobtailsrescue.org.uk

VIRAL HAEMORRHAGIC DISEASE (VHD)

HOW DOES A RABBIT GET VHD?

Direct contact with other rabbits, in water, in food, on people, clothing, objects, vermin, other domestic pets, bird droppings and through the air. It only takes 100 virus particles to make an infective dose (millions would fit on a pin head). The virus survives a long time in the environment so even indoor rabbits can be affected.

HOW CAN I STOP MY RABBIT GETTING VHD?

The only reliable way to protect your bunny is to have him/her vaccinated. The Nobivac Myxo-RHD vaccine protects them against both Myxomatosis and VHD classic strain 1. Please note for full protection your rabbit will also need an annual booster of Filavac -K C+V to protect them from the RHDV2 variant strain. These vaccinations should be given at least 14 days apart. Please contact your vet for advice.

WHAT HAPPENS IF MY RABBIT CATCHES VHD?

Some rabbits may drop dead without any outward signs of being ill. Others may suffer the following symptoms: high temperature, anorexia, lethargy, spasms, difficulty breathing, bleeding from the nose, mouth and anus. 80-100% of susceptible rabbits will die of the disease and very few recover even with the best care.

Every rabbit is at risk and the only way to protect your rabbit is to have them vaccinated and to keep their annual boosters up to date.



